

# Random runners are very lonely

Sebastian Czerwiński

**ABSTRACT.** Suppose that  $k$  runners having different constant speeds run laps on a circular track of unit length. The Lonely Runner Conjecture states that, sooner or later, any given runner will be at distance at least  $1/k$  from all the other runners. We prove that, with probability tending to one, a much stronger statement holds for random sets in which the bound  $1/k$  is replaced by  $1/2 - \varepsilon$ . The proof uses Fourier analytic methods. We also point out some consequences of our result for colouring of random integer distance graphs.

## 1. Introduction

Suppose that  $k$  runners run laps on a unit-length circular track. They all start together from the same point and run in the same direction with pairwise different constant speeds  $d_1, d_2, \dots, d_k$ . At a given time  $t$ , a runner is said to be *lonely* if no other runner is within a distance of  $1/k$ , both in front and rear. The Lonely Runner Conjecture states that for every runner there is a time at which he is lonely. For instance if  $k = 2$ , one can imagine easily that at some time or other, the two runners will find themselves on antipodal points of the circle, both becoming lonely at that moment.

To give a precise statement, let  $\mathbb{T} = [0, 1)$  denote the *circle* (the one-dimensional torus). For a real number  $x$ , let  $\{x\}$  be the fractional part of  $x$  (the position of  $x$  on the circle), and let  $\|x\|$  denote the distance of  $x$  to the nearest integer (the circular distance from  $\{x\}$  to zero). Notice that  $\|x - y\|$  is just the length of the shortest circular arc determined by the points  $\{x\}$  and  $\{y\}$  on the circle. It is not difficult to see that the following statement is equivalent to the Lonely Runner Conjecture.

**CONJECTURE 1.** *For every integer  $k \geq 1$  and for every set of positive integers  $\{d_1, d_2, \dots, d_k\}$  there exists a real number  $t$  such that*

$$\|td_i\| \geq \frac{1}{k+1}$$

*for all  $i = 1, 2, \dots, k$ .*

---

*Key words and phrases.* lonely runner problem, Diophantine approximation, integer distance graphs.

The above bound is sharp as is seen for the sets  $\{1, 2, \dots, k\}$ . The paper of Goddyn and Wong [11] contains items of interesting exemplars of such extremal sets. The problem was posed for the first time by Wills [16] in connection to Diophantine approximation. Cusick [8] raised the same question independently, as a view obstruction problem in Discrete Geometry (cf. [5]). Together with Pomerance [9], he confirmed the validity of the conjecture for  $k \leq 4$ . Bienia et al. [3] gave a simpler proof for  $k = 4$  and found interesting application to flows in graphs and matroids. Next the conjecture was proved for  $k = 5$  by Bohman et al. [4]. A simpler proof for that case was provided by Renault [13]. Recently the case  $k = 6$  was established by Barajas and Serra [2], using a new promising idea.

Let  $D = \{d_1, d_2, \dots, d_k\}$  be a set of  $k$  positive integers. Consider the quantity

$$\kappa(D) = \sup_{x \in \mathbb{T}} \min_{d_i \in D} \|xd_i\|$$

and the related function  $\kappa(k) = \inf \kappa(D)$ , where the infimum is taken over all  $k$ -element sets of positive integers. So, the Lonely Runner Conjecture states that  $\kappa(k) \geq \frac{1}{k+1}$ . The trivial bound is  $\kappa(k) \geq \frac{1}{2k}$ , as the sets  $\{x \in \mathbb{T} : \|xd_i\| < \frac{1}{2k}\}$  simply cannot cover the whole circle (since each of them is a union of  $d_i$  open arcs of length  $\frac{1}{kd_i}$  each). Surprisingly, nothing much better was proved so far. Currently the best general bound is

$$\kappa(k) \geq \frac{1}{2k - 1 + \frac{1}{2k-3}}$$

for every  $k \geq 5$  [6]. A slightly improved inequality  $\kappa(k) \geq \frac{1}{2k-3}$  holds when  $k \geq 4$  and  $2k - 3$  is prime [7]. Using the probabilistic argument we proved in [10] that every set  $D$  contains an element  $d$  such that

$$\kappa(D \setminus \{d\}) \geq \frac{1}{k}.$$

In this paper we prove another general result supporting the Lonely Runner Conjecture.

**THEOREM 1.** *Let  $k$  be a fixed positive integer and let  $\varepsilon > 0$  be fixed real number. Let  $D \subseteq \{1, 2, \dots, n\}$  be a  $k$ -element subset chosen uniformly at random. Then the probability that  $\kappa(D) \geq \frac{1}{2} - \varepsilon$  tends to 1 with  $n \rightarrow \infty$ .*

The proof uses elementary Fourier analytic technique for subsets of  $\mathbb{Z}_p$ . We give it in the next section. In the last section we point to a striking consequence of our result for colouring of integer distance graphs.

## 2. Proof of the main result

Let  $k$  be a fixed positive integer and let  $p \geq k$  be a prime number. For  $a \in \mathbb{Z}_p$ , let  $\|a\|_p = \min\{a, p - a\}$  be the circular distance from  $a$  to zero in  $\mathbb{Z}_p$ . We will need the following notion introduced by [14]. Let  $L$  be a fixed

positive integer. A set  $D = \{d_1, \dots, d_k\} \subseteq \mathbb{Z}_p$  is called  $L$ -independent in  $\mathbb{Z}_p$  if equation

$$d_1x_1 + d_2x_2 + \dots + d_kx_k = 0$$

has no solutions satisfying

$$0 < \sum_{i=1}^k \|x_i\|_p \leq L.$$

We will show that for appropriately chosen  $L$ , any  $L$ -independent set can be pushed away arbitrarily far from zero. Then we will demonstrate that for such  $L$ , almost every set in  $\mathbb{Z}_p$  is  $L$ -independent.

Let  $f : \mathbb{Z}_p \rightarrow \mathbb{C}$  be any function and let  $\hat{f} : \mathbb{Z}_p \rightarrow \mathbb{C}$  denote its Fourier transform, that is

$$\hat{f}(r) = \sum_{x \in \mathbb{Z}_p} f(x) \omega^{rx},$$

where  $\omega = e^{\frac{2\pi i}{p}}$ . For a set  $A \subseteq \mathbb{Z}_p$ , by  $A(x)$  we denote its characteristic function. We will make use of the following basic properties of the Fourier transform:

$$\textbf{(F1): } |\hat{f}(r)| = |\hat{f}(-r)| \text{ for every } r \in \mathbb{Z}_p.$$

$$\textbf{(F2): } f(x) = \frac{1}{p} \sum_{r \in \mathbb{Z}_p} \hat{f}(r) \omega^{-rx} \text{ for every } x \in \mathbb{Z}_p.$$

$$\textbf{(F3): } \hat{A}(0) = |A| \text{ for every subset of } \mathbb{Z}_p.$$

In the lemma below we give a bound for the Fourier coefficient  $\hat{A}(r)$  for the sets of the form

$$(*) \quad A = \{s, s+1, \dots, l\},$$

where  $l$  and  $s$  are elements of  $\mathbb{Z}_p$ , such that  $s < l$ . This bound does not depend on  $l$  and  $s$ . The following lemma can be easily proved, as for instance in [12](p. 39). We proved this for a reader convenience.

LEMMA 1. *If  $0 < r < \frac{p}{2}$ , then*

$$|\hat{A}(r)| \leq \frac{p}{2r}.$$

PROOF. By simple calculations we have

$$\begin{aligned} |\hat{A}(r)| &= \left| \sum_{x=s}^l \omega^{rx} \right| = \left| \frac{\omega^{r(l+1)} - \omega^{rs}}{\omega^r - 1} \right| = \\ &= \left| \frac{\omega^{\frac{r(l+s+1)}{2}}}{\omega^{\frac{r}{2}}} \cdot \frac{\omega^{\frac{r(l+1-s)}{2}} - \omega^{\frac{-r(l+1-s)}{2}}}{\omega^{\frac{r}{2}} - \omega^{\frac{-r}{2}}} \right| = \left| \frac{\sin(\frac{\pi r}{p})}{\sin(\frac{\pi r}{p})} \right|. \end{aligned}$$

Using inequality  $\sin(x) \geq \frac{2x}{\pi}$  for  $x < \frac{\pi}{2}$ , we get

$$\left| \hat{A}(r) \right| \leq \frac{p}{2r}. \quad \square$$

Now, we state and prove the aforementioned property of  $L$ -independent sets.

**THEOREM 2.** *Let  $0 < \varepsilon < \frac{1}{2}$  be a fixed real number. Let  $D$  be a  $k$ -element,  $L$ -independent set in  $\mathbb{Z}_p$ , where*

$$L > \sqrt{\frac{k^3 3^{k-1}}{2^{k+1} \varepsilon^{2k}}}.$$

*Then*

$$\kappa(D) \geq 1/2 - \varepsilon.$$

**PROOF.** Let

$$C = \{x \in \mathbb{Z}_p : (\frac{1}{4} - \frac{\varepsilon}{2})p < x < (\frac{1}{4} + \frac{\varepsilon}{2})p\}$$

and let  $C(x)$  be the characteristic function of the set  $C$ . Define convolution of two functions  $f$  and  $g$  by

$$(f * h)(x) = \sum_{y \in \mathbb{Z}_p} f(y) \cdot g(x - y).$$

Denote by  $B(x) = (C * C)(x)$  convolution of function  $C$  with itself. It is easy to see that  $\hat{B}(r) = \hat{C}(r) \cdot \hat{C}(r)$  for all  $r \in \mathbb{Z}_p$ .

So, if we find  $t \in \mathbb{Z}_p$  such that  $tD \subseteq \text{supp} B$ , where  $\text{supp} B = \{x \in \mathbb{Z}_p : B(x) \neq 0\}$ , then at the same time we push the set  $D$  away into the small arc  $(\frac{1}{2} - \varepsilon, \frac{1}{2} + \varepsilon)$  on the torus  $\mathbb{T}$ .

Then the expression

$$I = \sum_{t \in \mathbb{Z}_p} B(td_1)B(td_2) \cdots B(td_k)$$

counts those numbers  $t$  which push the set  $D$  away to a distance  $\frac{1}{2} - \varepsilon$  from zero. We will show that  $I \neq 0$ . From properties of the Fourier transform results that

$$I = \sum_{t \in \mathbb{Z}_p} \left( \frac{1}{p} \sum_{r_1 \in \mathbb{Z}_p} \hat{B}(r_1) \omega^{-td_1 r_1} \right) \cdots \left( \frac{1}{p} \sum_{r_k \in \mathbb{Z}_p} \hat{B}(r_k) \omega^{-td_k r_k} \right).$$

Denoting  $\vec{r} = (r_1, r_2, \dots, r_k)$ , we get

$$p^k I = \sum_{\vec{r} \in \mathbb{Z}_p^k} \hat{B}(r_1) \cdots \hat{B}(r_k) \sum_{t \in \mathbb{Z}_p} \omega^{-t(d_1 r_1 + \cdots + d_k r_k)}.$$

The expression  $\sum_t \omega^{-t(d_1 r_1 + \cdots + d_k r_k)}$  is equal to  $p$  when

$$(**) \quad d_1 r_1 + \cdots + d_k r_k \equiv 0 \pmod{p},$$

and is equal to zero in the contrary case. As a consequence we may write

$$p^{k-1}I = \sum_{\vec{r} \in \mathbb{Z}_p^k} \hat{B}(r_1) \cdots \hat{B}(r_k) R(\vec{r}),$$

where  $R(\vec{r}) = 1$  for  $r_1, \dots, r_k$  satisfying the equation (\*\*), and  $R(\vec{r}) = 0$  in the opposite situation. Since  $D$  is  $L$ -independent, the identity  $R(\vec{r}) = 1$  holds only for those  $r_1, \dots, r_k$  satisfying condition  $\sum_{i=1}^k \|r_i\|_p > L$ , or  $r_1 = r_2 = \dots = r_k = 0$ . Hence,

$$p^{k-1}I - |C|^{2k} = \sum_{\vec{r} \in \mathbb{Z}_p^k, \sum \|r_i\|_p > L} \hat{B}(r_1) \cdots \hat{B}(r_k),$$

as for  $r_i = 0$  the Fourier coefficient  $\hat{B}(r_i)$  is equal to square of the size of  $C$ . So, by showing that

$$|C|^{2k} > \sum_{\sum \|r_i\|_p > L} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_k) \right| R(\vec{r}),$$

we will confirm that  $I \neq 0$ .

The property of  $L$ -independence of the set  $D$  implies that in any non-trivial solution of (\*\*) there is some  $r_i$  satisfying  $\|r_i\|_p > \frac{L}{k}$ . The estimates for those  $r_i$

$$\left| \hat{B}(r_i) \right| = \left| \hat{C}(r_i) \right|^2 \leq \left( \frac{p}{2r_i} \right)^2 \leq \left( \frac{kp}{2L} \right)^2$$

results from Lemma 1.

Denote by  $\vec{r}_j = (r_1, \dots, r_{j-1}, r_{j+1}, \dots, r_k)$ , the vector  $\vec{r}$  with  $j^{th}$  coordinate missing. Substituting this to the previous sum we obtain

$$\begin{aligned} & \sum_{\sum \|r_i\|_p > L} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_k) \right| R(r_1, \dots, r_k) \\ & \leq \left( \frac{kp}{2L} \right)^2 \sum_{j=1}^k \sum_{\vec{r}_j \in \mathbb{Z}_p^{k-1}} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_{j-1}) \right| \left| \hat{B}(r_{j+1}) \right| \cdots \left| \hat{B}(r_k) \right| \\ & \leq k \left( \frac{kp}{2L} \right)^2 \sum_{\vec{r}_k \in \mathbb{Z}_p^{k-1}} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_{k-1}) \right|. \end{aligned}$$

The last sum may be estimated further. Let  $S_p = \{0, 1, \dots, \frac{p-1}{2}\}$  and we get

$$\begin{aligned} & \sum_{\vec{r}_k \in \mathbb{Z}_p^{k-1}} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_{k-1}) \right| \\ & \leq 2^{k-1} \sum_{\vec{r}_k \in S_p^{k-1}} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_{k-1}) \right|. \end{aligned}$$

Thus, applying Lemma 1 again we get

$$\begin{aligned} & \sum_{\sum \|r_i\|_p > L} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_k) \right| R(\vec{r}) \\ & \leq k \left( \frac{kp}{2L} \right)^2 \cdot 2^{k-1} \cdot \left( \frac{p^{k-1}}{2^{k-1}} \right)^2 \cdot \left( 1 + \sum_{r \in S_p} \frac{1}{r^2} \right)^{k-1} \\ & \leq k \left( \frac{kp}{2L} \right)^2 \cdot 2^{k-1} \cdot \left( \frac{p^{k-1}}{2^{k-1}} \right)^2 \cdot \left( 1 + \frac{\pi^2}{2} \right)^{k-1} \end{aligned}$$

since  $1 + \frac{\pi^2}{2} \leq 3$ , we obtain

$$\sum_{\sum \|r_i\|_p > L} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_k) \right| R(\vec{r}) \leq \frac{k^3 p^{2k} 3^{k-1}}{2^{k+1} L^2}.$$

So, by the assumption on  $L$  we obtain

$$\sum_{\sum \|r_i\|_p > L} \left| \hat{B}(r_1) \right| \cdots \left| \hat{B}(r_k) \right| R(\vec{r}) < (\varepsilon p)^{2k} \leq |C|^{2k}.$$

This completes the proof.  $\square$

PROOF OF THEOREM 1. Let  $L$  be a number satisfying inequalities

$$\sqrt{\frac{k^3 3^{k-1}}{2^{k+1} \varepsilon^{2k}}} < L < {}^{k+1}\sqrt{p}.$$

Such numbers  $L$  exist provided that  $p$  is sufficiently large. By Theorem 2,  $\kappa(D) \geq \frac{1}{2} - \varepsilon$  for every  $L$ -independent set  $D$ . We show that the second inequality implies that almost every set in  $\mathbb{Z}_p^*$  is  $L$ -independent. Indeed, the number of sets that are not  $L$ -independent is at most

$$(2L+1)^k \binom{p-1}{k-1}.$$

So, the fraction of those sets in  $\mathbb{Z}_p^*$  is equal

$$\frac{(2L+1)^k \binom{p-1}{k-1}}{\binom{p-1}{k}} = \frac{(2L+1)^k k}{p-k} < \frac{(2 {}^{k+1}\sqrt{p} + 1)^k k}{p-k}.$$

The last expression tends to zero with  $p$  tending to infinity. This completes the proof, as the ratios of two consecutive primes tend to one.  $\square$

### 3. Integer distance graphs

We conclude the paper with a remark concerning *integer distance graphs*. For a given set  $D$ , consider a graph  $G(D)$  whose vertices are positive integers, with two vertices  $a$  and  $b$  joined by an edge if and only if  $|a-b| \in D$ . Let  $\chi(D)$  denote the chromatic number of this graph. It is not hard to see that  $\chi(D) \leq |D| + 1$ .

To see a connection to parameter  $\kappa(D)$ , put  $N = \lceil \kappa(D)^{-1} \rceil$  and split the circle into  $N$  intervals  $I_i = [(i-1)/N, i/N)$ ,  $i = 1, 2, \dots, N$  (cf. [15]). Let

$t$  be a real number such that  $\min_{d \in D} \|dt\| = \kappa(D)$ . Then define a colouring  $c : \mathbb{N} \rightarrow \{1, 2, \dots, N\}$  by  $c(a) = i$  if and only if  $\{ta\} \in I_i$ . If  $c(a) = c(b)$  then  $\{ta\}$  and  $\{tb\}$  are in the same interval  $I_i$ . Hence  $\|ta - tb\| < 1/N \leq \kappa(D)$ , and therefore  $|a - b|$  is not in  $D$ . This means that  $c$  is a proper colouring of a graph  $G(D)$ . So, we have a relation

$$\chi(D) \leq \left\lceil \frac{1}{\kappa(D)} \right\rceil.$$

Now, by Theorem 1 we get that  $\chi(D) \leq 3$  for almost every graph  $G(D)$ .

A different proof of a stronger version of this result has been recently found by Alon [1]. He also extended the theorem for arbitrary Abelian groups, and posed many intriguing questions for general groups.

ACKNOWLEDGEMENT 1. *I would like to thank Tomasz Schoen for an inspiring idea of using independent sets, and to Jarek Grytczuk for stimulating discussions and help in preparation of the manuscript. I thank the anonymous referees for valuable suggestions concerning the merit of the paper. I also acknowledge a support from Polish Ministry of Science and Higher Education (MNiSW) (N N201 271335).*

## References

- [1] N. Alon, The chromatic number of random Cayley graphs, to appear.
- [2] J. Barajas, O. Serra, The lonely runner with seven runners, Electron. J. Combin. 15 (1) (2008), Paper R48.
- [3] W. Bienia, L. Goddyn, P. Gvozdzak, A. Sebö, M. Tarsi, Flows, view obstructions, and the lonely runner, J. Combin. Theory Ser. B 72 (1998), no. 1, 1–9.
- [4] T. Bohman, R. Holzman, D. Kleitman, Six lonely runners. In honor of Aviezri Fraenkel on the occasion of his 70th birthday, Electron. J. Combin. 8 (2001), no. 2, Research Paper 3, 49 pp. (electronic).
- [5] P. Brass, W. Moser, J. Pach, Research problems in discrete geometry, Springer, New York, 2005.
- [6] Y. G. Chen, View-obstruction problems and a generalization in  $E^n$ , Acta Math. Sinica, 37 (1994), 551–562.
- [7] Y. G. Chen and T. W. Cusick, The view-obstruction problem for  $n$ -dimensional cubes, J. Number Theory 74 (1999), 126–133.
- [8] T. W. Cusick, View-obstruction problems, Aequationes Math. 9 (1973), 165–170.
- [9] T. W. Cusick, C. Pomerance, View-obstruction problems. III, J. Number Theory 19 (1984), no. 2, 131–139.
- [10] S. Czerwiński, J. Grytczuk, Invisible runners in finite fields, Information Processing Letters 108 (2008), 64–67.
- [11] L. Goddyn, E. Wong, Tight instances of the Lonely Runner, INTEGERS: Elec. J. Comb. Num. Theory 6 (2006), #A38, 14pp.
- [12] H. L. Montgomery, Ten Lectures on the Interface between Analytic Number Theory and Harmonic Analysis, American Mathematical Society, 1994.
- [13] J. Renault, View-obstruction: a shorter proof for 6 lonely runners. Discrete Math, 287 (2004), no. 1-3, 93–101.
- [14] I. Ruzsa, Arithmetic progressions in sumset, Acta Arithmetica LX.2 (1991), 191–202.
- [15] I. Ruzsa, Z. Tuza, M. Voigt, Distance graphs with finite chromatic number, J. Combin. Theory, Ser. B, 85 (2002), 181–187.

- [16] J. M. Wills, Zwei Sätze über inhomogene diophantische Approximation von Irrationalzahlen, Monatsch. Math. 71 (1967), 263–269.
- [17] X. Zhu, Pattern periodic coloring of distance graphs, J. Combin. Theory Ser. B 73 (1998), 195–206.

FACULTY OF MATHEMATICS, COMPUTER SCIENCE, AND ECONOMETRICS, UNIVERSITY  
OF ZIELONA GÓRA, 65-546 ZIELONA GÓRA, POLAND  
*E-mail address:* s.czerwinski@wmie.uz.zgora.pl